OXFORD

INTERNATIONAL
AQA EXAMINATIONS

OXFORD AQA INTERNATIONAL AS

AS Physics

Data and formulae booklet

Insert

DATA - FUNDAMENTAL CONSTANTS AND VALUES

Quantity	Symbol	Value	Units
speed of light in vacuo	c	3.00×10^{8}	$\mathrm{m\ s^{-1}}$
permeability of free space	μ_0	$4\pi \times 10^{-7}$	$\mathrm{H\ m^{-1}}$
permittivity of free space	ε_0	8.85×10^{-12}	$\mathrm{F\ m^{-1}}$
magnitude of the charge of electron	e	1.60×10^{-19}	C
the Planck constant	h	6.63×10^{-34}	J s
gravitational constant	G	6.67×10^{-11}	$\mathrm{N\ m^2\ kg^{-2}}$
the Avogadro constant	N_A	6.02×10^{23}	$\mathrm{mol^{-1}}$
molar gas constant	R	8.31	$\mathrm{J\ K^{-1}\ mol^{-1}}$
the Boltzmann constant	k	1.38×10^{-23}	$\mathrm{J\ K^{-1}}$
the Stefan constant	σ	5.67×10^{-8}	$\mathrm{W\ m^{-2}\ K^{-4}}$
the Wien constant	α	2.90×10^{-3}	m K
electron rest mass (equivalent to 5.5×10^{-4} u)	m_e	9.11×10^{-31}	kg
electron charge/mass ratio	$\frac{e}{m_\mathrm{e}}$	1.76×10^{11}	$\mathrm{C\ kg^{-1}}$
proton rest mass (equivalent to 1.00728 u)	m_p	$1.67(3) \times 10^{-27}$	kg
proton charge/mass ratio	$\frac{e}{m_\mathrm{p}}$	9.58×10^{7}	$\mathrm{C\ kg^{-1}}$
neutron rest mass (equivalent to 1.00867 u)	m_n	$1.67(5) \times 10^{-27}$	kg
gravitational field strength	g	9.81	$\mathrm{N\ kg^{-1}}$
acceleration due to gravity	g	9.81	$\mathrm{m\ s^{-2}}$
atomic mass unit (1u is equivalent to 931.5 MeV)	u	1.661×10^{-27}	kg

ASTRONOMICAL DATA

Body	Mass/kg	Mean radius/m
Sun	1.99×10^{30}	6.96×10^{8}
Earth	5.97×10^{24}	6.37×10^{6}

GEOMETRICAL EQUATIONS

arc length	$= r\theta$
circumference of circle	$= 2\pi r$
area of circle	$= \pi r^2$
curved surface area of cylinder	$= 2\pi rh$
area of sphere	$= 4\pi r^2$
volume of sphere	$= \frac{4}{3}\pi r^3$

Unit 1

Mechanics and materials

moments $\text{moment} = Fd$

velocity and acceleration $v = \frac{\Delta s}{\Delta t}$ $a = \frac{\Delta v}{\Delta t}$

equations of motion $v = u + at$

$v^2 = u^2 + 2as$

$s = \left(\frac{u+v}{2}\right) t$

$s = ut + \frac{at^2}{2}$

force $F = ma$

$F = \frac{\Delta(mv)}{\Delta t}$

impulse $F\,\Delta t = \Delta(mv)$

work, energy and power $W = F\,s\cos\theta$

$E_\text{k} = \frac{1}{2}\,m\,v^2$ $\Delta E_\text{p} = mg\Delta h$

$P = \frac{\Delta W}{\Delta t}$, $P = Fv$

$efficiency = \frac{useful\ output\ power}{input\ power}$

density $\rho = \frac{m}{V}$

Hooke's law $F = k\,\Delta L$

Young modulus $= \frac{tensile\ stress}{tensile\ strain}$

tensile stress $= \frac{F}{A}$

tensile strain $= \frac{\Delta L}{L}$

energy stored $E = \frac{1}{2}F\Delta L$

Particles, radiation and radioactivity

inverse square law for γ radiation $I = \frac{I_0}{r^2}$

Unit 2

Electricity

current and pd $I = \frac{\Delta Q}{\Delta t}$ $V = \frac{W}{Q}$ $R = \frac{V}{I}$

resistivity $\rho = \frac{RA}{L}$

resistors in series $R_\text{T} = R_1 + R_2 + R_3 + \ldots$

resistors in parallel $\frac{1}{R_\text{T}} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \cdots$

energy transferred $E = IVt$

power $P = VI = I^2R = \frac{V^2}{R}$

emf $\varepsilon = \frac{E}{Q}$ $\varepsilon = I(R + r)$

Oscillations and waves

for a mass-spring system $T = 2\pi\sqrt{\frac{m}{k}}$

for a simple pendulum $T = 2\pi\sqrt{\frac{l}{g}}$

wave speed $c = f\lambda$ *period* $f = \frac{1}{T}$

first harmonic $f = \frac{1}{2l}\sqrt{\frac{T}{\mu}}$

fringe spacing $w = \frac{\lambda D}{s}$ *diffraction grating* $d\sin\theta = n\lambda$

refractive index of a substance s, $n = \frac{c}{c_\text{s}}$

for two different substances of refractive indices n_1 and n_2,

law of refraction $n_1\sin\theta_1 = n_2\sin\theta_2$

critical angle $\sin\theta_c = \frac{n_2}{n_1}$ for $n_1 > n_2$

photon energy $E = hf = \frac{hc}{\lambda}$

photoelectricity $hf = \phi + E_{\text{k (max)}}$

energy levels $hf = E_1 - E_2$

de Broglie Wavelength $\lambda = \frac{h}{p} = \frac{h}{mv}$

Copyright © 2017 Oxford International AQA Examinations and its licensors. All rights reserved.